

- b. The costs of manufacturing  $x_1$  dollars worth of B are given by the vector  $x_1\mathbf{b}$ , and the costs of manufacturing  $x_2$  dollars worth of C are given by  $x_2\mathbf{c}$ . Hence the total costs for both products are given by the vector  $x_1\mathbf{b} + x_2\mathbf{c}$ .  $\square$

## PRACTICE PROBLEMS

1. Prove that  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  for any  $\mathbf{u}$  and  $\mathbf{v}$  in  $\mathbb{R}^n$ .
2. For what value(s) of  $h$  will  $\mathbf{y}$  be in  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  if

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ -1 \\ -2 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 5 \\ -4 \\ -7 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} -3 \\ 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad \mathbf{y} = \begin{bmatrix} -4 \\ 3 \\ h \end{bmatrix}$$

## 1.3 EXERCISES

In Exercises 1 and 2, compute  $\mathbf{u} + \mathbf{v}$  and  $\mathbf{u} - 2\mathbf{v}$ .

1.  $\mathbf{u} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} -3 \\ -1 \end{bmatrix}$
2.  $\mathbf{u} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$

In Exercises 3 and 4, display the following vectors using arrows on an  $xy$ -graph:  $\mathbf{u}, \mathbf{v}, -\mathbf{v}, -2\mathbf{v}, \mathbf{u} + \mathbf{v}, \mathbf{u} - \mathbf{v}$ , and  $\mathbf{u} - 2\mathbf{v}$ . Notice that  $\mathbf{u} - \mathbf{v}$  is the vertex of a parallelogram whose other vertices are  $\mathbf{u}, 0$ , and  $-\mathbf{v}$ .


3.  $\mathbf{u}$  and  $\mathbf{v}$  as in Exercise 1
4.  $\mathbf{u}$  and  $\mathbf{v}$  as in Exercise 2

In Exercises 5 and 6, write a system of equations that is equivalent to the given vector equation.

5.  $x_1 \begin{bmatrix} 6 \\ -1 \\ 5 \end{bmatrix} + x_2 \begin{bmatrix} -3 \\ 4 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ -7 \\ -5 \end{bmatrix}$

6.  $x_1 \begin{bmatrix} -2 \\ 3 \end{bmatrix} + x_2 \begin{bmatrix} 8 \\ 5 \end{bmatrix} + x_3 \begin{bmatrix} 1 \\ -6 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

Use the accompanying figure to write each vector listed in Exercises 7 and 8 as a linear combination of  $\mathbf{u}$  and  $\mathbf{v}$ . Is every vector in  $\mathbb{R}^2$  a linear combination of  $\mathbf{u}$  and  $\mathbf{v}$ ?



7. Vectors  $\mathbf{a}, \mathbf{b}, \mathbf{c}$ , and  $\mathbf{d}$

8. Vectors  $\mathbf{w}, \mathbf{x}, \mathbf{y}$ , and  $\mathbf{z}$

In Exercises 9 and 10, write a vector equation that is equivalent to the given system of equations.

9.  $x_2 + 5x_3 = 0$   
 $4x_1 + 6x_2 - x_3 = 0$   
 $-x_1 + 3x_2 - 8x_3 = 0$
10.  $4x_1 + x_2 + 3x_3 = 9$   
 $x_1 - 7x_2 - 2x_3 = 2$   
 $8x_1 + 6x_2 - 5x_3 = 15$

In Exercises 11 and 12, determine if  $\mathbf{b}$  is a linear combination of  $\mathbf{a}_1, \mathbf{a}_2$ , and  $\mathbf{a}_3$ .

11.  $\mathbf{a}_1 = \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix}$ ,  $\mathbf{a}_2 = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}$ ,  $\mathbf{a}_3 = \begin{bmatrix} 5 \\ -6 \\ 8 \end{bmatrix}$ ,  $\mathbf{b} = \begin{bmatrix} 2 \\ -1 \\ 6 \end{bmatrix}$

12.  $\mathbf{a}_1 = \begin{bmatrix} 1 \\ -2 \\ 2 \end{bmatrix}$ ,  $\mathbf{a}_2 = \begin{bmatrix} 0 \\ 5 \\ 5 \end{bmatrix}$ ,  $\mathbf{a}_3 = \begin{bmatrix} 2 \\ 0 \\ 8 \end{bmatrix}$ ,  $\mathbf{b} = \begin{bmatrix} -5 \\ 11 \\ -7 \end{bmatrix}$

In Exercises 13 and 14, determine if  $\mathbf{b}$  is a linear combination of the vectors formed from the columns of the matrix  $A$ .

13.  $A = \begin{bmatrix} 1 & -4 & 2 \\ 0 & 3 & 5 \\ -2 & 8 & -4 \end{bmatrix}$ ,  $\mathbf{b} = \begin{bmatrix} 3 \\ -7 \\ -3 \end{bmatrix}$

14.  $A = \begin{bmatrix} 1 & -2 & -6 \\ 0 & 3 & 7 \\ 1 & -2 & 5 \end{bmatrix}$ ,  $\mathbf{b} = \begin{bmatrix} 11 \\ -5 \\ 9 \end{bmatrix}$

In Exercises 15 and 16, list five vectors in  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$ . For each vector, show the weights on  $\mathbf{v}_1$  and  $\mathbf{v}_2$  used to generate the vector and list the three entries of the vector. Do not make a sketch.

15.  $\mathbf{v}_1 = \begin{bmatrix} 7 \\ 1 \\ -6 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -5 \\ 3 \\ 0 \end{bmatrix}$

16.  $\mathbf{v}_1 = \begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -2 \\ 0 \\ 3 \end{bmatrix}$

17. Let  $\mathbf{a}_1 = \begin{bmatrix} 1 \\ 4 \\ -2 \end{bmatrix}$ ,  $\mathbf{a}_2 = \begin{bmatrix} -2 \\ -3 \\ 7 \end{bmatrix}$ , and  $\mathbf{b} = \begin{bmatrix} 4 \\ 1 \\ h \end{bmatrix}$ . For what value(s) of  $h$  is  $\mathbf{b}$  in the plane spanned by  $\mathbf{a}_1$  and  $\mathbf{a}_2$ ?

18. Let  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}$ ,  $\mathbf{v}_2 = \begin{bmatrix} -3 \\ 1 \\ 8 \end{bmatrix}$ , and  $\mathbf{y} = \begin{bmatrix} h \\ -5 \\ -3 \end{bmatrix}$ . For what value(s) of  $h$  is  $\mathbf{y}$  in the plane generated by  $\mathbf{v}_1$  and  $\mathbf{v}_2$ ?

19. Give a geometric description of  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$  for the vectors

$$\mathbf{v}_1 = \begin{bmatrix} 8 \\ 2 \\ -6 \end{bmatrix} \text{ and } \mathbf{v}_2 = \begin{bmatrix} 12 \\ 3 \\ -9 \end{bmatrix}$$

20. Give a geometric description of  $\text{Span}\{\mathbf{v}_1, \mathbf{v}_2\}$  for the vectors in Exercise 16.

21. Let  $\mathbf{u} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ . Show that  $\begin{bmatrix} h \\ k \end{bmatrix}$  is in  $\text{Span}\{\mathbf{u}, \mathbf{v}\}$  for all  $h$  and  $k$ .

22. Construct a  $3 \times 3$  matrix  $A$ , with nonzero entries, and a vector  $\mathbf{b}$  in  $\mathbb{R}^3$  such that  $\mathbf{b}$  is *not* in the set spanned by the columns of  $A$ .

In Exercises 23 and 24, mark each statement True or False. Justify each answer.

23. a. Another notation for the vector  $\begin{bmatrix} -4 \\ 3 \end{bmatrix}$  is  $[-4 \ 3]$ .

b. The points in the plane corresponding to  $\begin{bmatrix} -2 \\ 5 \end{bmatrix}$  and  $\begin{bmatrix} -5 \\ 2 \end{bmatrix}$  lie on a line through the origin.

c. An example of a linear combination of vectors  $\mathbf{v}_1$  and  $\mathbf{v}_2$  is the vector  $\frac{1}{2}\mathbf{v}_1$ .

d. The solution set of the linear system whose augmented matrix is  $[\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3 \ \mathbf{b}]$  is the same as the solution set of the equation  $x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + x_3\mathbf{a}_3 = \mathbf{b}$ .

e. The set  $\text{Span}\{\mathbf{u}, \mathbf{v}\}$  is always visualized as a plane through the origin.

24. a. Any list of five real numbers is a vector in  $\mathbb{R}^5$ .

b. The vector  $\mathbf{u}$  results when a vector  $\mathbf{u} - \mathbf{v}$  is added to the vector  $\mathbf{v}$ .

c. The weights  $c_1, \dots, c_p$  in a linear combination  $c_1\mathbf{v}_1 + \dots + c_p\mathbf{v}_p$  cannot all be zero.

d. When  $\mathbf{u}$  and  $\mathbf{v}$  are nonzero vectors,  $\text{Span}\{\mathbf{u}, \mathbf{v}\}$  contains the line through  $\mathbf{u}$  and the origin.

e. Asking whether the linear system corresponding to an augmented matrix  $[\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_3 \ \mathbf{b}]$  has a solution amounts to asking whether  $\mathbf{b}$  is in  $\text{Span}\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ .

25. Let  $A = \begin{bmatrix} 1 & 0 & -4 \\ 0 & 3 & -2 \\ -2 & 6 & 3 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} 4 \\ 1 \\ -4 \end{bmatrix}$ . Denote the

columns of  $A$  by  $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$ , and let  $W = \text{Span}\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ .

a. Is  $\mathbf{b}$  in  $\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ ? How many vectors are in  $\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3\}$ ?

b. Is  $\mathbf{b}$  in  $W$ ? How many vectors are in  $W$ ?

c. Show that  $\mathbf{a}_1$  is in  $W$ . [Hint: Row operations are unnecessary.]

26. Let  $A = \begin{bmatrix} 2 & 0 & 6 \\ -1 & 8 & 5 \\ 1 & -2 & 1 \end{bmatrix}$ , let  $\mathbf{b} = \begin{bmatrix} 10 \\ 3 \\ 3 \end{bmatrix}$ , and let  $W$  be the

set of all linear combinations of the columns of  $A$ .

a. Is  $\mathbf{b}$  in  $W$ ?

b. Show that the third column of  $A$  is in  $W$ .

27. A mining company has two mines. One day's operation at mine #1 produces ore that contains 20 metric tons of copper