

Chapter 2

Math 2890-001

Fall 2017

Due Oct 13

Name _____

Problems 1 – 13 are fair game for Exam 1.

1. (1 point) Let $A = \begin{pmatrix} 2 & 1 & 9 & 4 \\ 8 & 3 & 0 & 5 \\ 3 & 5 & 5 & 4 \end{pmatrix}$. Find A^T , the transpose of A .

answer: The transpose $A^T = \begin{pmatrix} 2 & 8 & 3 \\ 1 & 3 & 5 \\ 9 & 0 & 5 \\ 4 & 5 & 4 \end{pmatrix}$.

2. (1 point) Let $A = \begin{pmatrix} 9 & 2 & 4 \\ 2 & 8 & 9 \\ 4 & 1 & 3 \end{pmatrix}$. Is A symmetric?

answer: A is not symmetric since $A_{23} = 9 \neq 1 = A_{32}$.

3. (1 point) Let

$$A = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & -1 \\ 1 & -1 & 1 & 1 \\ 1 & -1 & -1 & 1 \\ 1 & 1 & -1 & -1 \end{pmatrix}.$$

Is A orthogonal? Explain your answer.

answer: A is not orthogonal since $I \neq A^T A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}.$

4. (3 points) For each of the following matrices, find the inverse if it exists. Show your work.

$$(a) A = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}.$$

$$\text{answer: } A^{-1} = \begin{pmatrix} 0 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 0 \end{pmatrix}.$$

$$(b) B = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}.$$

$$\text{answer: } A^{-1} = \begin{pmatrix} 1 & -2 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{pmatrix}.$$

$$(c) C = \begin{pmatrix} 3 & 2 & 1 \\ 2 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

$$\text{answer: } A^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & -2 \\ 1 & -2 & 1 \end{pmatrix}.$$

5. (1 point) Let $A = \begin{pmatrix} -2 & -1 \\ 8 & 8 \\ 0 & 1 \\ -1 & -8 \end{pmatrix}$ and $B = \begin{pmatrix} 3 & -2 \\ 6 & 5 \\ -6 & 4 \\ 8 & -3 \end{pmatrix}$.

Compute the sum $A + B$ (showing your work) if it is defined; otherwise, explain why it is not defined.

answer: The sum $A + B = \begin{pmatrix} 1 & -3 \\ 14 & 13 \\ -6 & 5 \\ 7 & -11 \end{pmatrix}$.

6. (1 point) Let $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}$ and $v = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}$.

Compute the product Av (showing your work) if it is defined; otherwise, explain why it is not defined.

answer: The product $Av = \begin{pmatrix} 10 \\ 4 \\ 1 \\ 6 \end{pmatrix}$.

7. (1 point) Let $v = (1 \ 2 \ 3 \ 4)$ and $A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \\ 1 & 1 & 1 \end{pmatrix}$.

Compute the product vA (showing your work) if it is defined; otherwise, explain why it is not defined.

answer: The product $vA = (5 \ 7 \ 14)$.

8. (1 point) Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 1 \\ 2 & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 2 & 1 & 1 & 3 \\ 1 & 2 & 3 & 1 \end{pmatrix}$.

Compute the product AB (showing your work) if it is defined; otherwise, explain why it is not defined.

answer: The product $AB = \begin{pmatrix} 4 & 5 & 7 & 5 \\ 7 & 5 & 6 & 10 \\ 6 & 6 & 8 & 8 \end{pmatrix}$.

9. (1 point) Suppose A is a 52×19 matrix and B is a 19×23 matrix. What size is the product AB if it is defined? Explain your answer.

10. (1 point) Let $A = \begin{pmatrix} 4 & 2 \\ 2 & 8 \\ 3 & 1 \\ 9 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 3 & 9 & 4 & 6 \\ 6 & 1 & 3 & 2 \end{pmatrix}$.

Compute column 3 of AB by first writing it as a linear combination of the columns of A . Show your work.

answer:

$$\begin{aligned} (AB)_{*3} &= A(B_{*3}) \\ &= \begin{pmatrix} 4 & 2 \\ 2 & 8 \\ 3 & 1 \\ 9 & 1 \end{pmatrix} \begin{pmatrix} 4 \\ 3 \end{pmatrix} \\ &= \begin{pmatrix} 4 \\ 2 \\ 3 \\ 9 \end{pmatrix} (4) + \begin{pmatrix} 2 \\ 8 \\ 1 \\ 1 \end{pmatrix} (3) \\ &= \begin{pmatrix} 22 \\ 32 \\ 15 \\ 39 \end{pmatrix} \end{aligned}$$

11. (1 point) Let $A = \begin{pmatrix} 4 & 2 \\ 2 & 8 \\ 3 & 1 \\ 9 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 3 & 9 & 4 & 6 \\ 6 & 1 & 3 & 2 \end{pmatrix}$.

Compute row 2 of AB by first writing it as a linear combination of the rows of B . Show your work.

answer:

$$\begin{aligned} (AB)_{2*} &= (A_{2*})B \\ &= (2 \ 8) \begin{pmatrix} 3 & 9 & 4 & 6 \\ 6 & 1 & 3 & 2 \end{pmatrix} \\ &= (2)(3 \ 9 \ 4 \ 6) + (8)(6 \ 1 \ 3 \ 2) \\ &= (54 \ 26 \ 32 \ 28) \end{aligned}$$

12. (1 point) Let $A = \begin{pmatrix} 4 & 2 \\ 2 & 8 \\ 3 & 1 \\ 9 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 3 & 9 & 4 & 6 \\ 6 & 1 & 3 & 2 \end{pmatrix}$.

Compute the entry in row 3, column 2 of the product AB . Show your work.

answer:

$$\begin{aligned} (AB)_{32} &= A_{3*}B_{*2} \\ &= (3 \ 1) \begin{pmatrix} 9 \\ 1 \end{pmatrix} \\ &= 28. \end{aligned}$$

13. (1 point) Let $A = \begin{pmatrix} 4 & 2 \\ 2 & 8 \\ 3 & 1 \\ 9 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 3 & 9 & 4 & 6 \\ 6 & 1 & 3 & 2 \end{pmatrix}$.

Write the product AB as a sum of 2 rank one matrices. Show your work.

answer:

$$\begin{aligned} AB &= A_{*1}B_{1*} + A_{*2}B_{2*} \\ &= \begin{pmatrix} 4 \\ 2 \\ 3 \\ 9 \end{pmatrix} (3 \ 9 \ 4 \ 6) + \begin{pmatrix} 2 \\ 8 \\ 1 \\ 1 \end{pmatrix} (6 \ 1 \ 3 \ 2) \\ &= \begin{pmatrix} 12 & 36 & 16 & 24 \\ 6 & 18 & 8 & 12 \\ 9 & 27 & 12 & 18 \\ 17 & 81 & 36 & 54 \end{pmatrix} + \begin{pmatrix} 12 & 2 & 6 & 4 \\ 48 & 8 & 24 & 16 \\ 6 & 1 & 3 & 2 \\ 6 & 1 & 3 & 2 \end{pmatrix} \end{aligned}$$

The remaining problems will not be covered on Exam 1.

14. (1 point) Let

$$A = \begin{pmatrix} 5 & -4 & -1 \\ -10 & 4 & 2 \\ 25 & -8 & -3 \end{pmatrix} \quad L = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 5 & -3 & 1 \end{pmatrix}$$
$$U = \begin{pmatrix} 5 & -4 & -1 \\ 0 & -4 & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} -10 \\ -8 \\ 48 \end{pmatrix}.$$

Use the LU factorization $A = LU$ to solve the matrix equation $Ax = b$. Show your work.

answer: The vector $y = \begin{pmatrix} -10 \\ -28 \\ 14 \end{pmatrix}$ is the solution of the system $Ly = b$.

and the vector $x = \begin{pmatrix} 5 \\ 7 \\ 7 \end{pmatrix}$ is the solution of $Ux = y$ and so of $Ax = b$.

15. (1 point) Let

$$A = \begin{pmatrix} 4 & -2 & 5 \\ 12 & -3 & 18 \\ -4 & -13 & -17 \end{pmatrix}.$$

Use Wedderburn rank reduction (or Gaussian Elimination) to find the LDU (or LU) factorization of the matrix A . Show your work.

answer: Either $A = LDU$ where

$$L = \begin{pmatrix} 4 & 0 & 0 \\ 12 & 3 & 0 \\ -4 & -15 & 3 \end{pmatrix} \quad D = \begin{pmatrix} 1/4 & 0 & 0 \\ 0 & 1/3 & 0 \\ 0 & 0 & 1/3 \end{pmatrix} \quad U = \begin{pmatrix} 4 & -2 & 5 \\ 0 & 3 & 3 \\ 0 & 0 & 3 \end{pmatrix},$$

or $A = LU$ where

$$L = \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ -1 & -5 & 1 \end{pmatrix} \quad U = \begin{pmatrix} 4 & -2 & 5 \\ 0 & 3 & 3 \\ 0 & 0 & 3 \end{pmatrix}.$$

16. (1 point) Let

$$A = \begin{pmatrix} 15 & 0 & 13 \\ -20 & 8 & -4 \\ 10 & -13 & -1 \end{pmatrix}.$$

Use Wedderburn rank reduction (or Gaussian Elimination with Partial Pivoting) to find the permuted LDU (or permuted LU) factorization of the matrix A . Show your work.

answer: Either $A = LDU$ where

$$L = \begin{pmatrix} 15 & 6 & 8 \\ -20 & 0 & 0 \\ 10 & -9 & 0 \end{pmatrix} \quad D = \begin{pmatrix} -1/20 & 0 & 0 \\ 0 & -1/9 & 0 \\ 0 & 0 & 1/8 \end{pmatrix} \quad U = \begin{pmatrix} -20 & 8 & -4 \\ 0 & -9 & -3 \\ 0 & 0 & 8 \end{pmatrix},$$

or $A = LU$ where

$$L = \begin{pmatrix} -0.75 & -0.6667 & 1 \\ 1 & 0 & 0 \\ -0.5 & 1 & 0 \end{pmatrix} \quad U = \begin{pmatrix} -20 & 8 & -4 \\ 0 & -9 & -3 \\ 0 & 0 & 8 \end{pmatrix},$$

17. (2 points) Let

$$A = \begin{pmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ 1 & 2 & 1 \end{pmatrix} \quad \text{and} \quad v = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}.$$

- (a) Is v in the column space of A ? Show your work.
(b) Is v in the null space of A ? Show your work.

answer: (a) Since $(A|v)$ does **not** have a pivot in the last column $Ax = v$ is consistent, and v **is** in the column space of A .

(b) v **is** in the null space of A since $Av = 0$.

18. (2 points) Let

$$A = \begin{pmatrix} 1 & 2 & 0 \\ 2 & 5 & 1 \\ 1 & 1 & -1 \end{pmatrix}.$$

- (a) Find a nonzero vector in the column space of A . Show your work.
(b) Find a nonzero vector in the null space of A . Show your work.

answer: (a) I found the vector $\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ in the column space. You'll probably find a different vector.

(b) I found the vector $\begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$ in the null space. You'll probably find a different vector.

19. (1 point) Let

$$A = \begin{pmatrix} -5 & 5 & 2 & 25 & 18 & -2 \\ -2 & 0 & 2 & 10 & 2 & -1 \\ -2 & -4 & 4 & 8 & -8 & -4 \\ -5 & -3 & 5 & 16 & -1 & 0 \\ 2 & 2 & 0 & 6 & 0 & -2 \end{pmatrix}.$$

Find bases for $\text{Col}(A)$ and $\text{Nul}(A)$.

Hint: The reduced row echelon form of A is

$$\begin{pmatrix} 1 & 0 & 0 & 0 & -2 & 0 \\ 0 & 1 & 0 & 3 & 2 & 0 \\ 0 & 0 & 1 & 5 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

answer: The columns of the matrix $\begin{pmatrix} -5 & 5 & 2 & -2 \\ -2 & 0 & 2 & -1 \\ -2 & -4 & 4 & -4 \\ -5 & -3 & 5 & 0 \\ 2 & 2 & 0 & -2 \end{pmatrix}$ are a basis for the column space of A .

The columns of the matrix $\begin{pmatrix} 0 & 2 \\ -3 & -2 \\ -5 & 1 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$ are a basis for the null space of A .

20. (1 point) Let

$$A = \begin{pmatrix} 58 & -15 & -144 & -204 & 11 \\ 101 & -27 & -249 & -357 & -4 \\ -45 & 12 & 111 & 159 & 1 \\ 15 & -4 & -37 & -53 & 0 \end{pmatrix}.$$

Find bases for $\text{Col}(A)$, $\text{Nul}(A)$, $\text{Col}(A^T)$ and $\text{Nul}(A^T)$.

Hint: If you form the matrix $(A|I)$ and use row operations to put the A part in reduced row echelon form you get $(R|S)$ where

$$R = \begin{pmatrix} 1 & 0 & -3 & -3 & 0 \\ 0 & 1 & -2 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad S = \begin{pmatrix} 1 & 3 & 1 & -21 \\ 4 & 13 & 8 & -79 \\ -1 & -7 & -16 & 3 \\ 1 & 7 & 17 & 0 \end{pmatrix}.$$

answer: The columns of the matrix $\begin{pmatrix} 58 & -15 & 11 \\ 101 & -27 & -4 \\ -45 & 12 & 1 \\ 15 & -4 & 0 \end{pmatrix}$ form a basis for the column space of A .

The columns of the matrix $\begin{pmatrix} 3 & 3 \\ 2 & -2 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$ form a basis for the null space of A .

The columns of the matrix $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -3 & -2 & 0 \\ -3 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ form a basis for the column space of A^T .

The columns of the matrix $\begin{pmatrix} 1 \\ 7 \\ 17 \\ 0 \end{pmatrix}$ form a basis for the null space of A^T .

21. (1 point) Let

$$A = \begin{pmatrix} 1 & 2 & 8 & 1 & 5 \\ 2 & 1 & 7 & 2 & 7 \\ 1 & 1 & 5 & 0 & 2 \\ 3 & 2 & 12 & 1 & 7 \end{pmatrix}.$$

Find the rank of the matrix A . Explain your answer.

answer: The reduced row echelon form of A is

$$\begin{pmatrix} 1 & 0 & 2 & 0 & 1 \\ 0 & 1 & 3 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

a matrix having 3 pivots, so the rank of A is 3.