

### 13.3 Conditional Probability and Intersection of Events



1

**Conditional probability** is the probability of one event (F) happening assuming that another event (E) does.

Examples:

- probability that someone is happy given that they just won \$\$\$.
- probability that someone passes an exam given that they did not study.

The probability that F happens given that E does is denoted  $P(F|E)$

It is read "probability of F given E"

Example: Flip 2 coins for an experiment.

What is the probability that a Head is flipped given that the 1<sup>st</sup> coin was a Tail?

3

Example: Flip 2 coins for an experiment.

What is the probability that a Head is flipped given that the 1<sup>st</sup> coin was a Tail?

The event we assume happened was that the 1<sup>st</sup> was a Tail.

The event among those is that there is a Head

4

Example: Flip 2 coins for an experiment.

What is the probability that a Head is flipped given that the 1<sup>st</sup> coin was a Tail?

The event we assume happened was that the 1<sup>st</sup> was a Tail.  
 $\{ TH, TT \}$

The event among those is that there is a Head.  
 $\{ TH \}$

$$P(H | 1^{\text{st}} \text{ is } T) = 1/2$$

5

Note that for the full experiment there are 4 outcomes, but we are only interested when the "given" outcome occurs.

6

Example: Roll a die for an experiment.

What is the probability it is odd given that the value was a prime number?

7

Example: Roll a die for an experiment.

What is the probability it is odd given that the value was a prime number?

The event assumed to happen was that the value was prime.

Among those the event is when is it odd.

8

Example: Roll a die for an experiment.

What is the probability it is odd given that the value was a prime number?

The event assumed to happen was that the value was prime.  
 $\{2, 3, 5\}$

Among those the event is when is it odd.  
 $\{3, 5\}$

$P(\text{odd} \mid \text{prime}) = 2/3$

9

The previous examples lead to a way to count  $P(F \mid E)$  by a formula:

**SPECIAL RULE FOR COMPUTING  $P(F \mid E)$  BY COUNTING** If  $E$  and  $F$  are events in a sample space with equally likely outcomes, then  $P(F \mid E) = \frac{n(E \cap F)}{n(E)}$ .

Recall:  $E \cap F = E \text{ intersect } F = E \text{ and } F$

10

Example: Two dice are rolled (order matters)

What is the probability that 1<sup>st</sup> die is 3 given that the sum is 4?

11

Example: Two dice are rolled (order matters)

What is the probability that 1<sup>st</sup> die is 3 given that the sum is 4?

Event "sum is 4"

Event "sum is 4 and 1<sup>st</sup> die is 3"

12

Example: Two dice are rolled (order matters)

What is the probability that 1<sup>st</sup> die is 3 given that the sum is 4?

Event "sum is 4"  
 $\{ (1, 3), (2, 2), (3, 1) \}$

Event "sum is 4 and 1<sup>st</sup> die is 3"  
 $\{ (3, 1) \}$

$$P(1^{\text{st}} \text{ is } 3 \mid \text{sum is } 4) = \frac{n(1^{\text{st}} \text{ is } 3 \text{ and sum is } 4)}{n(\text{sum is } 4)} = 1/3$$

## Conditional Probability

- Example: Assume that we roll two dice and the total showing is greater than nine. What is the probability that the total is odd?

11/01/13  
 © 2010 Pearson Education, Inc. All rights reserved.

(continued on next slide)  
 Section 14.3, Slide 14

## Conditional Probability

- Example: Assume that we roll two dice and the total showing is greater than nine. What is the probability that the total is odd?


- Solution: This sample space has 36 equally likely outcomes. We will let  $G$  be the event "we roll a total greater than nine" and let  $O$  be the event "the total is odd." Therefore,  
 $G = \{(4, 6), (5, 5), (5, 6), (6, 4), (6, 5), (6, 6)\}$ .

11/01/13  
 © 2010 Pearson Education, Inc. All rights reserved.

(continued on next slide)  
 Section 14.3, Slide 15

## Conditional Probability

We now seek all pairs that give an odd total – the diagram below shows that there are two.



$$P(O|G) = \frac{n(O \cap G)}{n(G)} = \frac{2}{6} = \frac{1}{3}$$

11/01/13  
 © 2010 Pearson Education, Inc. All rights reserved.

Section 14.3, Slide 16

Note that  $P(E \mid F)$  and  $P(F \mid E)$  are different.

Example:

If  $n(E) = 4$ ,  $n(F) = 8$ , and  
 $n(E \cap F) = 2$

$$P(E \mid F) = P(E \cap F) / P(F) = 2/4 = 1/2$$

$$P(F \mid E) = P(E \cap F) / P(E) = 2/4 = 1/2$$

17

By multiplying through on the formula...

**RULE FOR COMPUTING THE PROBABILITY OF THE INTERSECTION OF EVENTS** If  $E$  and  $F$  are two events, then

$$P(E \cap F) = P(E) \cdot P(F \mid E).$$

18

In testing for a disease, a test works 90% of the time given that the person has the disease.  
10% of the people have the disease.

What is the probability that someone has the disease and the test works?

19

In testing for a disease, a test works 90% of the time given that the person has the disease.  
10% of the people have the disease.

What is the probability that someone has the disease and the test works?

$$P(\text{test works} \mid \text{disease}) = 0.9$$

$$P(\text{disease}) = 0.1$$

$$\begin{aligned} P(\text{test works and disease}) &= P(\text{test works} \mid \text{disease}) \times P(\text{disease}) \\ &= 0.9 \times 0.1 \\ &= 0.09 \end{aligned}$$

## The Intersection of Events

- Example: Assume your professor has written questions on 10 assigned readings on cards and you are to randomly select two cards and write an essay on them. If you have read 8 of the 10 readings, what is your probability of getting two questions that you can answer?

11/01/13

© 2010 Pearson Education, Inc. All rights reserved.

(continued on next slide)

Section 14.3, Slide 21

## The Intersection of Events

- Example: Assume your professor has written questions on 10 assigned readings on cards and you are to randomly select two cards and write an essay on them. If you have read 8 of the 10 readings, what is your probability of getting two questions that you can answer?

- Solution: Let  $A$  be “you can answer the first question;” and  $B$  be “you can answer the second question.”

11/01/13

© 2010 Pearson Education, Inc. All rights reserved.

(continued on next slide)

Section 14.3, Slide 22

## The Intersection of Events

We need to calculate

probability you can answer the first question      probability you can answer the second question, given that you answered the first question

$$P(A \cap B) = P(A) \cdot P(B \mid A).$$

We may compute the following probabilities:

$$P(A) = \frac{8}{10} \quad P(B \mid A) = \frac{7}{9}$$

$$P(A \cap B) = P(A) \cdot P(B \mid A) = \frac{8}{10} \cdot \frac{7}{9} = \frac{56}{90} \approx 0.62$$

11/01/13

© 2010 Pearson Education, Inc. All rights reserved.

23

Section 14.3, Slide 23